

Indications of the possible observation of the lowest-lying 1^{-+} QCD state

Kwei-Chou Yang¹

¹*Department of Physics, Chung-Yuan Christian University, Chung-Li, Taiwan 320, R.O.C.*

(Dated: March, 17, 2007)

We discuss properties of 1^{-+} exotic meson within the framework of the QCD field-theoretic approach. We estimate the mass of the lowest-lying 1^{-+} exotic meson using renormalization-improved QCD sum rules, and find that the mass lies around 1.26 ± 0.15 GeV, in good agreement with the $\pi_1(1400)$ data. This state should be expected in QCD. We find that the mass for the lowest-lying strange 1^{-+} meson is 1.31 ± 0.19 GeV. Our result hints that the $K^*(1410)$ may be the lowest-lying 1^{-+} nonet state.

PACS numbers: 12.39.Mk, 11.55.Hx, 12.38.Aw, 12.38.Lg

I. INTRODUCTION

Quantum Chromodynamics (QCD) is the presently accepted theory of strong interactions among quarks and gluons. Perturbative QCD has been developed in great detail and tested successfully. However, the physics of the nonperturbative QCD seems to be a much more difficult task. The rapid development of the lattice QCD theory may offer some answers to the nonperturbative dynamics.

In addition to normal mesons ($\bar{q}q$) and baryons (qqq), which can be built up from the naive quark model, QCD allows the existence of exotic states which can be glueballs, hybrid mesons (the bound states of $\bar{q}qg$), tetraquark mesons (the bound states of $\bar{q}q\bar{q}q$), and other multi-parton states. These exotic states, which go beyond the description of the naive quark model, can offer the direct evidence concerning confining properties of QCD.

In the past years possible evidences have accumulated for the existence of 1^{-+} exotic states [1]. Now we have two 1^{-+} exotic states below 2 GeV. A negatively charged exotic state, $\pi_1(1400)$, with $J^{PC} = 1^{-+}$ was observed in $\pi^- p \rightarrow \eta \pi^- p$ [2, 3] and in $\bar{p} n \rightarrow \pi^0 \pi^- \eta$ [4]. The corresponding neutral state was reported by the Crystal Barrel [5] and E852 [6] collaborations in the reactions of $\bar{p} p \rightarrow \pi^0 \pi^0 \eta$ and $\pi^- p \rightarrow \eta \pi^0 n$, respectively, where the decay channel $\eta \pi$ is isovector and hence cannot be confused with a glueball. Unlike the charged $\eta \pi^-$ channel, the charge conjugate is a good quantum number in the neutral $\eta \pi^0$ system, where the η was detected in its 2γ decay mode [6] and very recently in its $\pi^+ \pi^- \pi^0$ decay mode in the E852 experiment [7]. The advantage of detecting the $\eta \rightarrow \pi^+ \pi^- \pi^0$ mode over the all-neutral final state is that the decaying vertex is determined by charged tracks. The mass of the neutral exotic 1^{-+} state observed in the very recent E852 experiment is $1257 \pm 20 \pm 25$ MeV [7], which is lower than the mass 1360 ± 25 MeV in the Crystal Barrel measurement. The world average is 1376 ± 19 MeV [1].

Another observed 1^{-+} exotic state, $\pi_1(1600)$, has also attracted much attention. In contrast to $\pi_1(1400)$, the η'/π coupling of this state is unexpectedly stronger than

the $\eta \pi$ coupling although the phase space favors the latter. Due to the above property, it was argued that the $\pi_1(1400)$ may be favored for a four-quark state, while the $\pi_1(1600)$ may be a hybrid meson [8].

Theoretically, the compositions of observed 1^{-+} exotic states remain unclear. The flux-tube model [8] and lattice calculations [9] predict the lowest-lying 1^{-+} hybrid meson to have a mass of about 1.9 GeV. Moreover, the former predicts that the 1^{-+} hybrid meson are dominated by $b_1 \pi$ and $f_1 \pi$ decays in contrast with the $\pi_1(1600)$ experimental results where the three final states $b_1 \pi$, $\eta' \pi$ and $\rho \pi$ are of comparable strength, $1 : 1.0 \pm 0.3 : 1.5 \pm 0.5$ [10].

To evaluate the mass of the lowest-lying 1^{-+} hybrid meson in the QCD sum rule approach, in the literature, people used

$$J_\mu(x) = i \bar{d}(x) \gamma_\alpha g_s G_\mu^\alpha(x) u(x),$$

as the relevant interpolating current in the calculation of the two-point correlation function. The earliest QCD sum rule calculations were done in Refs. [11, 12], where the radiative corrections and anomalous dimensions of J_μ and some operators in operator-product-expansion (OPE) series were not known. To obtain more precise estimate, the radiative corrections, which are sizable, were then computed in Refs. [13, 14]. Unfortunately, the mass sum rule result is highly sensitive to the parameter s_0 which models the threshold of the excited states, and does not show s_0 stability. Thus, Jin and Körner conservatively estimate that the mass of the lowest-lying 1^{-+} hybrid meson is larger than 1.55 GeV [14]. Nevertheless, Chetyrkin and Narison argue that s_0 may lie between 3.5 GeV² and 4.5 GeV², so that the mass of the lowest-lying 1^{-+} hybrid meson is about 1.6 ~ 1.7 GeV [13].

To clarify the inconsistencies in the literature, in the present work, we study the mass of the lowest-lying 1^{-+} hybrid meson using the QCD sum rule technique. Instead of J_μ , we will use

$$J(x) = n^\beta n^\mu \bar{d}(x) \sigma_{\alpha\beta} g_s G_\mu^\alpha(x) u(x).$$

Here and in what follows, n^μ is a light-like vector, which satisfies $n^2 = 0$. So far no people adopt the above current to study the mass of the so-called hybrid meson. The

reason that we use J is because the residue for J coupling to the 1^{-+} hybrid meson determines the normalization of the twist-3 light-cone distribution amplitude (LCDA) of the lowest-lying 1^{-+} meson, while, for J_μ , the residue corresponds to the normalization of twist-4 LCDA. For a relevant reaction, the amplitude related to J_μ is relatively suppressed by m/Q , as compared with that due to J , where Q corresponds to the scale of the reaction and m is the mass of the 1^{-+} meson. Thus J may be more suitable to use to study the lowest-lying 1^{-+} meson than J_μ .

Note that the renormalization(RG)-improvement is always necessary in the method of QCD sum rules because the hadronic mass is determined by the maximum stability of the sum rule within the Borel window, where the Borel mass is actually the renormalization scale. Unlike the vector current $\bar{q}\gamma_\mu q$ which satisfies the current conservation and is scale-independent, J as well as J_μ is a scale-dependent operator, similar to the baryonic cases [15, 16], and its anomalous dimension may result in significant improvement in the QCD sum rule result.

The rest of this paper is organized as follows. In Sec. II, we make a brief description of LCDAs for the 1^{-+} exotic meson in language of the QCD field theory. The LCDAs play an essential rôle in the QCD description of hard exclusive processes. Using the QCD sum rule technique and adopting the current J , in Sec. III we are devoted to the calculation of the mass sum rule of the lowest-lying 1^{-+} meson. To clarify the discrepancy between our result and those given in the literature, we revisit the previous QCD sum rule studies in Sec. IV, where the current J_μ is adopted. Finally, Sec. V contains the conclusions and discussions.

II. THE FRAMEWORK OF QCD

Unlike the model-building approach, a 1^{-+} exotic meson in language of the QCD field theory is described in terms of a set of Fock states for which each state has the same quantum number as the exotic meson:

$$|1^{-+}\rangle = \psi_{q\bar{q}}|\bar{q}q\rangle + \psi_{q\bar{q}g}|q\bar{q}g\rangle + \psi_{q\bar{q}q\bar{q}}|q\bar{q}q\bar{q}\rangle + \dots, \quad (1)$$

where ψ_i are distribution amplitudes and the dots denote the higher Fock states. In contrast to the usual intuition of people, actually a nonlocal $\psi_{q\bar{q}}$ does not vanish and is antisymmetric under interchange of momentum fractions of $\bar{q}$ and q in SU(3) limit. For instance, projecting a lowest-lying 1^{-+} meson (π_1) along the light-cone ($z^2 = 0$), the leading-twist LCDAs $\phi_{\parallel,\perp}$ are defined as

$$\begin{aligned} \langle 0|\bar{q}_1(z) \not{z} q_2(0)|\pi_1(P, \lambda)\rangle &= m_{\pi_1}^2 \epsilon^{(\lambda)} \cdot z [\phi_{\parallel}], \\ \langle 0|\bar{q}_1(z) \sigma_{\perp\nu} z^\nu q_2(0)|\pi_1(P, \lambda)\rangle &= im_{\pi_1} \epsilon_{\perp}^{(\lambda)} p \cdot z [\phi_{\perp}], \end{aligned} \quad (2)$$

where the notation $[\phi_{\parallel,\perp}] \equiv \int_0^1 du e^{-i\bar{u}pz} \phi_{\parallel,\perp}(u)$ and $p_\mu = P_\mu - z_\mu m_{\pi_1}^2/(2pz)$ are introduced with $\bar{u}(u)$ being

the momentum fraction carried by $\bar{q}_1(q_2)$, and the non-local quark-antiquark pair, connected by the Wilson line which is not shown, is at light-like separation. Considering the G-parity in Eq. (2), it can be known that $\phi_{\parallel,\perp}$ are antisymmetric under interchange $u \leftrightarrow \bar{u}$ in SU(3) limit, i.e., the amplitudes vanish in the $z \rightarrow 0$ limit (but *not* in the $z^2 \rightarrow 0$ limit). This property was first used in the study of the deep exclusive electro-production involving a lowest-lying 1^{-+} exotic meson [17]. In analogy to the leading LCDAs, all twist-three two-parton LCDAs for a 1^{-+} exotic meson are also antisymmetric under interchange $u \leftrightarrow \bar{u}$ in SU(3) limit due to the G-parity.

$\psi_{q\bar{q}g}$ can be non-vanishing under interchange of momentum fractions of quarks. Similar to the case of vector mesons [18], using non-local 3-parton gauge-invariant operators to project amplitudes of the $|q\bar{q}g\rangle$, we have three twist-3 3-parton LCDAs for a 1^{-+} exotic meson, built up by a quark, an antiquark, and a gluon, where two of LCDAs are symmetric under interchange of momentum fractions of the quark and antiquark in the SU(3) limit, while one is antisymmetric.

III. EVALUATION OF THE MASS OF THE LOWEST-LYING 1^{-+} MESON

Adopting the local gauge-invariant current ¹

$$J(x) = n^\beta n^\mu \bar{d}(x) \sigma_{\alpha\beta} g_s G_\mu^\alpha(x) u(x), \quad (3)$$

we shall employ the QCD sum rules [19] to evaluate the mass of the lowest-lying 1^{-+} exotic meson. $J(x)$ is G-parity odd, the same as the 1^{-+} isovector state. The residue of J coupled to the 1^{-+} state is defined as

$$\langle 0|J(0)|1^{-+}(p, \lambda)\rangle = f_{3,1^{-+}}^\perp m_{1^{-+}} (\epsilon^{(\lambda)} \cdot n)(p \cdot n). \quad (4)$$

The residue constant determines the normalization of one of twist-3 3-parton LCDAs, and is also the coefficient with conformal spin 7/2 in the conformal partial wave expansion for the LCDA [20].

The method of QCD sum rule approaches the bound state problem in QCD from the perturbative region, where non-perturbative quantities, such as some condensates, may contribute significant corrections in the OPE series. Through the study of the relevant correlation function and the idea of the quark-hadron duality, the corresponding hadronic properties, like masses, decay constants, form factors, etc., can be thus obtained.

We consider the two-point correlation function

$$i \int d^4x e^{iqx} \langle 0|T J(x) J^\dagger(0)|0\rangle = \Pi(q^2)(q \cdot n)^4. \quad (5)$$

¹ The other possible choice of the twist-3 current is to consider $n^\beta n^\alpha \bar{d}(x) g_s \gamma_\beta G_{\mu\alpha}(x) u(x)$. One can also consider the 4-quark operator relevant to the $|q\bar{q}q\bar{q}\rangle$ Fock state; however the resulting sum rule will be clouded by the factorization of condensates.

It should be noted that $J(x)$ can couple not only to 1^{-+} states but also to 0^{++} states as

$$\langle 0|J(0)|0^{++}(p, \lambda)\rangle = -2f_{3,S}(p \cdot n)^2, \quad (6)$$

where the lowest-lying state in the 0^{++} channel is $a_0(980)$. Therefore, to extract the lowest-lying meson corresponding to the 1^{-+} channel, at the hadron level of Eq. (5) we shall consider two lowest-lying states. Our final result will indicate that one of the two lowest-lying states is $a_0(980)$ and the other one is the lowest-lying 1^{-+} meson, i.e., the mass of the latter is lower than that of the $a_0(1450)$, the first excited state in the 0^{++} channel. We approximate the correlation function as

$$\frac{4(f_{3,a_0})^2}{m_{a_0}^2 - q^2} + \frac{(f_{3,\pi_1}^\perp)^2}{m_{\pi_1}^2 - q^2} = \frac{1}{\pi} \int_0^{s_0} ds \frac{\text{Im}\Pi^{\text{OPE}}}{s - q^2}, \quad (7)$$

where Π^{OPE} is the OPE result at the quark-gluon level, and s_0 is the threshold of higher resonances. We apply the Borel transformation to both sides of Eq. (7) to improve the convergence of the OPE series and to suppress contributions from higher resonances. The sum rule for the first two lowest-lying states can be written as

$$4e^{-m_{a_0}^2/M^2} f_{3,a_0}^2 + e^{-m_{\pi_1}^2/M^2} (f_{3,\pi_1}^\perp)^2 = \frac{1}{\pi} \int_0^{s_0} ds e^{-s/M^2} \text{Im}\Pi^{\text{OPE}}(s), \quad (8)$$

where M is the so-called Borel mass. We have checked that our Π^{OPE} result is consistent with that given in Ref. [21], where the authors combine Π^{OPE} with another correlation function result for calculating the twist-3 parameter of pseudoscalar mesons. Note that the authors of Ref. [21] did not perform the sum rule analysis as we do here. To improve further the M^2 dependence of the various terms using the RG equation. Thus we need to multiply each OPE term by a coefficient $L^{(-2\gamma_J + \gamma_n)/2b}$, where $L \equiv \alpha_s(M)/\alpha_s(\mu)$, $\gamma_J = 2(7C_F/3 + N_c)$ is the anomalous dimension of J , γ_n is the anomalous dimension of the corresponding operator, and $b = (11N_c - 2n_f)/3$ with N_c and n_f being numbers of colors and flavors, respectively [22]. The anomalous dimensions of operators appearing in the OPE series can be found in Ref. [16]. The full RG-improved sum rule thus reads

$$\begin{aligned} & 4e^{-m_{a_0}^2/M^2} [f_{3,a_0}(\mu)]^2 + e^{-m_{\pi_1}^2/M^2} [f_{3,\pi_1}^\perp(\mu)]^2 \\ &= \left\{ \frac{\alpha_s(\mu)}{360\pi^3} \int_0^{s_0} s e^{-s/M^2} ds + \frac{89}{5184} \frac{\alpha_s(\mu)}{\pi^2} \langle \alpha_s G^2 \rangle_\mu \right. \\ &+ \frac{\alpha_s(\mu)}{18\pi} (m_u \langle \bar{u}u \rangle_\mu + m_d \langle \bar{d}d \rangle_\mu) \\ &- \frac{\alpha_s(\mu)}{108\pi} \frac{1}{M^2} (m_u \langle \bar{u}g_s \sigma G u \rangle_\mu + m_d \langle \bar{d}g_s \sigma G d \rangle_\mu) L^{14/3b} \\ &+ \frac{71}{729} \frac{[\alpha_s(\mu)]^2}{M^2} (\langle \bar{u}u \rangle_\mu^2 + \langle \bar{d}d \rangle_\mu^2) L^{(b-s)/b} \\ &\left. - \frac{32}{81} \frac{[\alpha_s(\mu)]^2}{M^2} \langle \bar{u}u \rangle_\mu \langle \bar{d}d \rangle_\mu L^{(b-s)/b} \right\} L^{(b-\gamma_J)/b}. \quad (9) \end{aligned}$$

To subtract the contribution arising from the lowest-lying scalar meson, we further evaluate the following non-diagonal correlation function

$$i \int d^4x e^{iqx} \langle 0|TJ(x)\bar{u}(0)d(0)|0\rangle = \bar{\Pi}(q^2)(q \cdot n)^2, \quad (10)$$

where the scalar current can couple to the lowest-lying scalar meson $a_0(980)$:

$$\langle 0|\bar{u}(0)d(0)|a_0(980)\rangle = m_{a_0} f_{a_0}. \quad (11)$$

A similar non-diagonal correlation function was carried out in Ref. [21], where there are additional γ_5 's in two currents. Similarly, we thus obtain

$$\begin{aligned} & 2e^{-m_{a_0}^2/M^2} f_{3,a_0}(\mu) f_{a_0}(\mu) m_{a_0} \\ &= \left\{ \frac{\alpha_s(\mu)}{72\pi^3} L^{-1} \int_0^{s_0} s e^{-s/M^2} ds + \frac{1}{12\pi} \langle \alpha_s G^2 \rangle_\mu L^{-1} \right. \\ &- \frac{\alpha_s(\mu)}{9\pi} (m_u \langle \bar{u}u \rangle_\mu + m_d \langle \bar{d}d \rangle_\mu) L^{-1} \\ &+ \frac{2\alpha_s(\mu)}{9\pi} (m_u \langle \bar{d}d \rangle_\mu + m_d \langle \bar{u}u \rangle_\mu) L^{-1} \\ &\times \left[\frac{8}{3} + \gamma_E + \ln \frac{\mu^2}{M^2} - \text{Ei}\left(-\frac{s_0}{M^2}\right) \right] \\ &+ \frac{1}{6M^2} (m_u \langle \bar{d}g_s \sigma G d \rangle_\mu + m_d \langle \bar{u}g_s \sigma G u \rangle_\mu) L^{(14-3b)/3b} \\ &+ \frac{16}{27} \frac{\pi \alpha_s}{M^2} (\langle \bar{u}u \rangle_\mu^2 + \langle \bar{d}d \rangle_\mu^2) L^{-8/b} \\ &\left. - \frac{16}{9} \frac{\pi \alpha_s}{M^2} \langle \bar{u}u \rangle_\mu \langle \bar{d}d \rangle_\mu L^{-8/b} \right\} L^{(2b-\gamma_J)/2b}. \quad (12) \end{aligned}$$

Note that the RG-improvement for QCD sum rule results is very important. For instance, $L^{(2b-\gamma_J)/2b} = 1$ (or $L^{(b-\gamma_J)/b} = 1$) at $M^2 = 1 \text{ GeV}^2$, while the value becomes 0.93 (or 1.08) at $M^2 = 2 \text{ GeV}^2$. In calculating the mass sum rule for the lowest-lying 1^{-+} exotic state, in Eq. (9) we substitute f_{3,a_0} with the expression given in Eq. (12), where use of $f_{a_0} = (0.380 \pm 0.015) \text{ GeV}$ and $\bar{s}_0 = (3.0 \pm 0.2) \text{ GeV}^2$ have been made [23]. In the numerical analysis, we shall use the following values at the scale $\mu = 1 \text{ GeV}$:

$$\begin{aligned} & \langle \alpha_s G_{\mu\nu}^a G^{a\mu\nu} \rangle = (0.474 \pm 0.120) \text{ GeV}^4/(4\pi), \\ & \langle \bar{u}u \rangle \cong \langle \bar{d}d \rangle = -(0.24 \pm 0.01)^3 \text{ GeV}^3, \\ & \langle \bar{u}g_s \sigma G u \rangle \cong \langle \bar{d}g_s \sigma G d \rangle = (0.8 \pm 0.1) \langle \bar{u}u \rangle, \\ & (m_u + m_d)/2 = (5 \pm 2) \text{ MeV}. \end{aligned} \quad (13)$$

Consequently, the mass sum rule for the lowest-lying 1^{-+} resonance can be obtained by taking the logarithm of both sides of Eq. (9) and then applying the differential operator $M^4 \partial/\partial M^2$ to them. We choose the the Borel window $1 \text{ GeV}^2 < M^2 < 2 \text{ GeV}^2$, where the contribution originating from higher resonances (and the continuum), which is defined as

$$\frac{\frac{1}{\pi} \int_{s_0}^{\infty} ds e^{-s/M^2} \text{Im}\Pi^{\text{OPE}}(s)}{\frac{1}{\pi} \int_0^{\infty} ds e^{-s/M^2} \text{Im}\Pi^{\text{OPE}}(s)}, \quad (14)$$

is less than 53% and the highest OPE term at the quark-gluon level is no more than 13%. Both are well under control. To obtain a reliable estimate for the mass sum rule result, the contributions arising from higher resonances and the highest OPE term at the quark-gluon level cannot be too large. The excited threshold s_0 can be determined when the most stable plateau of the mass sum rule result is obtained within the Borel window. Note that no stable plateau can be obtained for the mass sum rule if the sum rule result does not contain the RG-corrections. (See also the discussions in Sec. IV.)

Numerically, we get the mass for the lowest-lying 1^{-+} exotic meson:

$$m_{\pi_1} = (1.26 \pm 0.15) \text{ GeV}, \quad (15)$$

corresponding to $s_0 = 2.5 \pm 0.7 \text{ GeV}^2$. The result for m_{π_1} versus M^2 is shown in Fig. 1, where the mass is very stable within the window. From Eq. (15), we know that the second lowest-lying meson for states coupling to the operator J should not be $a_0(1450)$ since the resulting mass is lower than the experimental result of $m_{a_0(1450)} = (1.474 \pm 0.019) \text{ GeV}$ [1]. It should be stressed that the procedure for performing the RG-improvement on the “mass” sum rule is very important. If the anomalous dimension of the current J was neglected, the stable sum rule could not be obtained within the Borel window and the resulting mass was reduced by 300 MeV.

We further study the existence of the 1^{-+} nonet. In evaluating the mass for the lowest-lying strange 1^{-+} exotic meson, we use $m_s(1 \text{ GeV}) = (135 \pm 15) \text{ MeV}$, $\langle \bar{s}g_s\sigma Gs \rangle / \langle \bar{u}u \rangle \approx \langle \bar{s}g_s\sigma Gs \rangle / \langle \bar{u}u \rangle = 0.85 \pm 0.05$ as additional inputs. Because it is still questionable whether the $\kappa(800)$ exists, we therefore consider the following two possible scenarios. In scenario 1, the $\kappa(800)$ is treated as the lowest-lying strange scalar meson with mass being $0.8 \pm 0.1 \text{ GeV}$ and $f_\kappa = 0.37 \pm 0.02 \text{ GeV}$, which corresponds to $\bar{s}_0 = 2.9 \pm 0.2 \text{ GeV}^2$, while in scenario 2, the $K_0^*(1430)$ is considered as the lowest-lying strange scalar meson with $f_{K_0^*(1430)} = 0.37 \pm 0.02 \text{ GeV}$, which corresponds to $\bar{s}_0 = 3.6 \pm 0.3 \text{ GeV}^2$. (The values are updated from Ref. [23].) The results are depicted in Fig. 2. Using an arbitrary set of allowed inputs, within the Borel window, the mass is stable only for scenario 2, which hints that the κ may not be a real particle or suitable in the sum rule study due to its large width. Because it is not stable in scenario 1, we assume $s_0 = 2.6 \pm 0.7 \text{ GeV}^2$, consistent with the case of π_1 . The result in scenario 2 is

$$m_{K^*(1^{-+})} = 1.31 \pm 0.19 \text{ GeV}, \quad (16)$$

corresponding to $s_0 = 2.3 \pm 0.9 \text{ GeV}^2$.

IV. THE MASS SUM RULE, AS DERIVED FROM THE CORRELATION FUNCTION GIVEN IN THE LITERATURE, REVISITED

Before concluding this paper, to clarify why our result given in the previous section differs from those given in




FIG. 1: The mass for the lowest-lying 1^{-+} exotic meson as a function of the Borel mass squared M^2 . The solid curve is obtained by using the central values of input parameters. The region between two dashed lines is variation of the mass within the allowed range of input parameters.




FIG. 2: The same as Fig. 1 but for the mass of the lowest-lying strange 1^{-+} exotic meson. The light (blue) and heavy (red) curves are for scenario 1 and 2, respectively, corresponding to $s_0 = 2.7 \pm 0.7$ and $2.3 \pm 0.9 \text{ GeV}^2$.

the literature, where a larger mass ($\gtrsim 1.55 \text{ GeV}$) for the lowest-lying 1^{-+} hybrid meson was obtained, we take into account the following correction function

$$i \int d^4x e^{iqx} \langle 0 | J_\mu(x) J_\nu^\dagger(0) | 0 \rangle = (q_\mu q_\nu - q^2 g_{\mu\nu}) \Pi_v(q^2) + q_\mu q_\nu \Pi_s(q^2), \quad (17)$$

with

$$J_\mu(x) = i \bar{d}(x) \gamma_\alpha g_s G_\mu^\alpha(x) u(x), \quad (18)$$

which was adopted in Refs. [13, 14]. The resulting sum rule can be written in the following form

$$e^{-m_{\pi_1}^2/M^2} m_{\pi_1}^4 (f_{4,\pi_1})^2 = \frac{1}{\pi} \int_0^{s_0} ds e^{-s/M^2} \text{Im} \Pi_v^{\text{OPE}}(s), \quad (19)$$

where

$$\langle 0 | J_\mu(0) | \pi_1(p, \lambda) \rangle = f_{4,\pi_1} m_{\pi_1}^2 \varepsilon_\mu^{(\lambda)}. \quad (20)$$

The f_{4,π_1} determines the normalization of twist-4 LCDAs of the lowest-lying 1^{-+} meson. The detailed OPE result for Π_v can be found in Refs. [13, 14], where the radiative corrections have been calculated.

The mass can then be obtained by applying $(M^4 \partial / \partial M^2 \ln)$ to both sides of Eq. (19). Before proceeding, three remarks are in order. First, although the anomalous dimension of J_μ was computed in Ref. [13], such a correction was not taken into account in Refs. [13, 14]. The scale dependence of the operator J_μ is given by $J_\mu(M) = J_\mu(\mu) L^{32/(9b)}$ [13]. Second, the scale-dependence of the strong-coupling constant is ignored in Refs. [13, 14]. Third, for theoretical completeness, the RG effect should be included in the mass sum rule (19) as done in Eq. (9), so that the right hand side of Eq. (19) should be multiplied by the overall factor $L^{1-64/(9b)}$. Such a factor equal to 1 at $M^2 = 1 \text{ GeV}^2$, but becomes ~ 0.95 at $M^2 = 2 \text{ GeV}^2$.

In Fig. 3, we plot the mass sum rule as a function of the Borel mass squared within $1 \text{ GeV}^2 < M^2 < 2 \text{ GeV}^2$. The light curves (the upper set) are the results,² where we simply fix the scale at 2 GeV for input parameters and use $s_0 = 4 \text{ GeV}^2$, which consist with that obtained by Jin and Körner [14] (see Fig. 3 in Ref. [14]). Moreover, if RG effects are not considered, as in Ref. [14], there is no plateau for any value of s_0 . Nevertheless, the stable plateau can be reached for a typical value of s_0 if we take into account the RG corrections in the sum rule. See the heavy curves (the lower set) in Fig. 3. The corresponding results are

$$\begin{aligned} m_{\pi_1} &= (1.18 \pm 0.09) \text{ GeV}, \\ s_0 &= (2.4_{-0.2}^{+0.4}) \text{ GeV}^2, \end{aligned} \quad (21)$$

where the errors are due to variation of the input parameters. Note that the plateau covers the range for $M^2 \sim 1.5 \text{ GeV}^2$. Unfortunately, for $1 \text{ GeV}^2 < M^2 < 2 \text{ GeV}^2$ and $s_0 = 2.4 \text{ GeV}^2$, the contribution originating from higher resonances (and the continuum) lies between 72% and 95%, which is too large, so that the resulting mass is less reliable.

V. CONCLUSIONS AND DISCUSSIONS

In this work we have studied the mass of the lowest-lying 1^{-+} hybrid meson using the QCD sum rule technique. Instead of $J_\mu(x) = id(x)\gamma_\alpha g_s G_\mu^\alpha(x)u(x)$ which is always adopted in the literature, we use, for the first

time,

$$J(x) = z^\beta z^\mu \bar{d}(x) \sigma_{\alpha\beta} g_s G_\mu^\alpha(x) u(x),$$

as the interpolating current of the lowest-lying 1^{-+} hybrid meson. Our main results are as follows:

- Using J as the interpolating current of the 1^{-+} exotic mesons, which are the so-called 1^{-+} hybrid mesons (see the discussions in Sec. II), we have obtained the mass for the lowest-lying 1^{-+} exotic meson to be $(1.26 \pm 0.15) \text{ GeV}$. The threshold of excited states, s_0 , as well as the resulting mass can be thus determined due to the existence of the most stable plateau of the mass sum rule result within the Borel window, where the contributions arising from the resonances and the highest OPE term at the quark-gluon level are well under control. The plateau for the mass result versus the Borel mass squared exists only when the RG-corrections are taken into account in the sum rule. Our result is in good agreement with the world average of data for the mass of $\pi_1(1400)$, which is $m_{\pi_1(1400)} = (1.376 \pm 0.019) \text{ GeV}$.
- On the other hand, to clarify the discrepancy between our result and those given in the literature for which J_μ is used as the interpolating field of the 1^{-+} hybrid meson, we have reexamined the previous sum rule calculation. (See Sec. IV.) As in Ref. [14], the curve for the resulting mass versus the Borel mass squared has no plateau region for any value of s_0 if RG corrections are not considered. Unfortunately, with RG-corrections, the plateau covers the range for $M^2 \sim 1.5 \text{ GeV}^2$, where the contribution arising from higher resonances is $\gtrsim 80\%$ in the sum rule result, so that the resulting mass is less reliable.




FIG. 3: The same as Fig. 1 but from the sum rule given in Eq. (19). The heavy (red) curves are obtained from RG-improved sum rule, where the solid curve corresponds to $s_0 = 2.4 \text{ GeV}^2$. The light (blue) curves, using $s_0 = 4.0 \text{ GeV}^2$ and the scale $\mu = 2 \text{ GeV}$ for α_s and remaining parameters, do not contain RG corrections.

² In Ref. [13], Chetyrkin and Narison use the finite energy sum rule approach in the analysis and argue that s_0 may lie between 3.5 GeV^2 and 4.5 GeV^2 . They obtain the mass of the lowest-lying 1^{-+} hybrid meson to be $1.6 \sim 1.7 \text{ GeV}$. In their analysis, the used parameter $1/\tau$ roughly agrees with $M^2 \approx (1.6 \sim 2) \text{ GeV}^2$. We thus find that the light curves in Fig. 3 are consistent with the result obtained by Chetyrkin and Narison.

- Using J in the sum rule study, we also obtain the mass for the strange 1^{-+} hybrid meson: $m_{K^*(1^{-+})} = 1.31 \pm 0.19$ GeV. In the Particle Data Book (PDG) [1], the two strange mesons, $K^*(1410)$ and $K^*(1680)$, are currently assigned to be 2^3S_1 and 1^3D_1 states, respectively. However, because the $K^*(1410)$ is too light as compared with the remaining 2^3S_1 nonet states, therefore it could be replaced by $K^*(1680)$ as the 2^3S_1 state. If so, our result hints that the $K^*(1410)$ is very likely to belong to the lowest-lying 1^{-+} nonet.
- We have presented a discussion for properties of the 1^{-+} exotic meson (which is usually called the 1^{-+} hybrid meson) based on the QCD field theory. Although $q\bar{q}g$ Fock states of the exotic meson are suppressed (see also discussions in Sec. I), the twist-2 distribution amplitudes, which are antisymmetric under interchange of the quark and antiquark momentum fractions in the SU(3) limit, give main contributions to the deep exclusive electroproduction involving a lowest-lying 1^{-+} exotic meson. Thus, from the point of view of QCD, the cross section of the above reaction is still sizable and the exotic (hybrid) meson can be studied in

experiments at JLAB, HERMES or Compass [17]. Furthermore, because of the non-small first Gegenbauer moment of the twist-2 distribution amplitudes, the branching ratios for two-body B decays involving the $\pi_1(1400)$ [24] could be of order 10^{-6} , which are easily accessible at the B factories.

In conclusion, we may have evidence for the existence of 1^{-+} exotic mesons. These states are allowed in Quantum Chromodynamics but not in the conventional quark model. Theorists tried to predict these states, especially with mass $\lesssim 1.6$ GeV, but did not succeed. In this paper, we have shown that the $\pi_1(1400)$ state is indeed expected in QCD. We also predict the mass of the lowest-lying strange 1^{-+} exotic meson which is a little larger than $m_{\pi_1(1400)}$. Thus, the observation of the $\pi_1(1400)$ state can be the direct evidence of QCD in explaining the confinement mechanism of strong interactions.

Acknowledgments

This work was supported in part by the National Science Council of R.O.C. under Grant No: NSC95-2112-M-033-001.

-
- [1] Particle Data Group, W. M. Yao *et al.*, J. Phys. G **33**, 1 (2006).
 - [2] D.R. Thompson *et al.*, Phys. Rev. Lett. **79**, 1630 (1997).
 - [3] S.U. Chung *et al.*, Phys. Rev. D **60**, 092001 (1999).
 - [4] A. Abele *et al.*, Phys. Lett. B **423**, 175 (1998).
 - [5] A. Abele *et al.*, Phys. Lett. B **446**, 349 (1999).
 - [6] A.R. Dzierba *et al.*, Phys. Rev. D **67**, 094015 (2003).
 - [7] G. Adams [E852 Collaboration], arXiv:hep-ex/0612062.
 - [8] F.E. Close and H.J. Lipkin, Phys. Lett. B **196**, 245 (1987); F.E. Close and P.R. Page, Nucl. Phys. B **443**, 233 (1995).
 - [9] C.W. Bernard *et al.*, Phys. Rev. D **56**, 7039 (1997).
 - [10] C. Amsler and N.A. Tornqvist, Phys. Rept. **389**, 61 (2004).
 - [11] J. Govaerts, F. de Viron, D. Gusbin and J. Weyers, Nucl. Phys. B **248**, 1 (1984).
 - [12] I.I. Balitsky, D. Diakonov and A.V. Yung, Z. Phys. C **33** (1986) 265.
 - [13] K.G. Chetyrkin and S. Narison, Phys. Lett. B **485**, 145 (2000).
 - [14] H.Y. Jin and J.G. Korner, Phys. Rev. D **64**, 074002 (2001).
 - [15] B. L. Ioffe, Nucl. Phys. B **188**, 317 (1981) [Erratum-ibid. B **191**, 591 (1981)].
 - [16] K. C. Yang, W. Y. P. Hwang, E. M. Henley and L. S. Kisslinger, Phys. Rev. D **47** (1993) 3001.
 - [17] I. V. Anikin, B. Pire, L. Szymanowski, O. V. Teryaev and S. Wallon, Phys. Rev. D **70**, 011501(R) (2004); I.V. Anikin, B. Pire, L. Szymanowski, O. V. Teryaev and S. Wallon, Phys. Rev. D **71**, 034021 (2005).
 - [18] P. Ball, V.M. Braun, Y. Koike and K. Tanaka, Nucl. Phys. B **529**, 323 (1998).
 - [19] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, Nucl. Phys. B **147** (1979) 385.
 - [20] V.M. Braun, G.P. Korchemsky and D. Muller, Prog. Part. Nucl. Phys. **51**, 311 (2003).
 - [21] P. Ball, V.M. Braun and A. Lenz, JHEP **0605**, 004 (2006).
 - [22] K. C. Yang, Nucl. Phys. B **776**, 187 (2007) [arXiv:0705.0692 [hep-ph]].
 - [23] H.Y. Cheng, C.K. Chua and K.C. Yang, Phys. Rev. D **73**, 014017 (2006).
 - [24] K.C. Yang, in preparation.